

**This Page Is Inserted by IFW Operations
and is not a part of the Official Record**

BEST AVAILABLE IMAGES

**Defective images within this document are accurate representations of
~~the original documents submitted by the applicant.~~**

Defects in the images may include (but are not limited to):

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/33, 31/56	A1	(11) International Publication Number: WO 99/25333 (43) International Publication Date: 27 May 1999 (27.05.99)
(21) International Application Number: PCT/US98/23386 (22) International Filing Date: 3 November 1998 (03.11.98) (30) Priority Data: 60/066,198 19 November 1997 (19.11.97) US (71) Applicant (for all designated States except US): HUMANETICS CORPORATION [US/US]; Interchange Tower, Suite 1205, 600 South Highway, St. Louis Park, MN 55426-1205 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): LARDY, Henry, A. [US/US]; 1829 Thorstrand Road, Madison, WI 53705 (US). WEEKS, Charles, E. [US/US]; 5276 Lakeview Avenue, White Bear Lake, MN 55110 (US). (74) Agent: SHERRILL, Michael, S.; Michael S. Sherrill Law Offices, Suite 200, 4756 Banning Avenue, White Bear Lake, MN 55110-3205 (US).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: USE OF $\Delta 5$ -ANDROSTENE- 3β -OL-7,17-DIONE IN THE TREATMENT OF LUPUS ERYTHEMATOSUS (57) Abstract Lupus erythematosus can be treated by administering therapeutic amounts of $\Delta 5$ -androstene- 3β -ol-7,17-dione and metabolizable precursors thereof, such as $\Delta 5$ -androstene- 3β -acetoxy-7,17-dione, which are readily metabolized <i>in vivo</i> to $\Delta 5$ -androstene- 3β -ol-7,17-dione but are not appreciably metabolizable <i>in vivo</i> to androgens, estrogens or dehydroepiandrosterone. Such treatment can be prophylactic, ameliorative or curative in nature.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**USE OF Δ^5 -ANDROSTENE-3 β -OL-7,17-DIONE
IN THE TREATMENT OF
LUPUS ERYTHEMATOSUS**

5

FIELD OF THE INVENTION

This invention broadly relates to treatment strategies for lupus erythematosus. More specifically, the invention relates to prophylactic, ameliorative and curative drug therapies for lupus erythematosus.

10

BACKGROUND

Lupus erythematosus is an autoimmune disorder which may, but does not always, affect many different organ systems in an affected individual. Lupus erythematosus (hereinafter "lupus") may affect the heart, lungs, skin, joints, kidneys, nervous system, lymph gland system, blood cells and/or blood vessels. Certain forms of lupus affect only or predominantly the skin. These forms of lupus are the most visible manifestations of the disease.

20

The immune system of the body is a complex and elaborate mechanism of protection from foreign substances. The immune system provides resistance to foreign cells and substances (*e.g.*, bacteria or virus) that may cause injury, as well as searching out abnormal cells (*e.g.*, cancerous cells) within the body for destruction. The invading or abnormal cells are neutralized by the immune system humoral and cellular components, including lymphocytes, antibodies, mediating (regulating such as lymphokines) systems, and effector (cytotoxic) cells. As an example, the immune system can recognize the invading or abnormal cells (antigens) within the body and produce antibodies (proteins) which attach to the recognized antigens, leading to their removal. Autoimmunity occurs when the immune system produces antibodies to normal cells in the body. This produces inflammation of normal tissue, resulting in

25
30

damage and loss of function. In other instances, antibodies attach to antigens within the blood plasma to form immune complexes that may be deposited in normal tissue resulting in inflammation and damage.

5 Women are more susceptible to lupus than men. Over 90% of lupus patients are females aged 13-40 years. Laboratory tests for the presence of lupus include the LE Cell Test, the Anti-Nuclear Antibody Test, and the test for Anti-DNA-Antibodies. Lupus is, however, often recognized by particular clinical manifestations including: (i) arthritis (occurring in 90-95% of persons with systemic lupus), (ii) skin changes, such as a
10 photosensitive induced "butterfly" rash across the bridge of the nose, across the cheeks and/or beneath the eyes, and/or red, raised and scaly patches, known as discoid lupus, anywhere on the body (occurring in 75-80% of persons with lupus), (iii) hematologic abnormalities, such as anemia, leukopenia, and thrombocytopenia (occurring in about 50% of persons with lupus), (iv) kidney impairment (occurring in about 50% of persons
15 with lupus), (v) heart or lung disease, such as an irritation of the heart or lung lining causing pericarditis or pleurisy (occurring in about 30% of persons with lupus), and (vi) neuropsychiatric changes (occurring in about 10% to 20% of persons with lupus).

Etiology

20

Lupus erythematosus can be divided into subsets which may or may not have overlapping characteristics: discoid lupus, subacute cutaneous lupus, drug-induced lupus and systemic lupus. Patients in whom the disease seems to be confined to the skin are differentiated from those with systemic or "disseminated" involvement.

25

Discoid lupus, also called chronic discoid lupus or chronic cutaneous lupus produces lesions over the face, but sometimes spread more extensively across the body. The lesions are usually well circumscribed, disk-like plaques of scaling erythema, tending to clear centrally with scarring, depigmentation and atrophy. Photosensitivity is
30 a common feature but may be absent. Roughly 80% to 90% of patients with discoid lupus lesions will not develop any signs and symptoms of systemic lupus.

Subacute cutaneous lupus produces wide spread skin lesions over the trunk and extremities of the patient. The lesions are apt to be less discrete than those of discoid lupus, more widespread, with temporary depigmentation and telangiectasia without scarring or atrophy. Loss of hair without scarring is common, and mild systemic disease, especially involving the joints, accompanied by fever and malaise are often present.

Another distinct subset of lupus is that associated with reaction to certain drugs, appropriately known as drug induced lupus. Serious organ involvement is rare and prognosis for this subset of lupus is excellent, provided the disease is recognized and the offending medication discontinued.

Systemic lupus is a chronic autoimmune disease that often has a relapsing course. The primary therapeutic approach for systemic lupus is to achieve and maintain adequate suppression of the disease with minimal drug mediated side effects. Evaluation of specific symptoms and clinical findings establish the type and extent of organ involvement and overall disease activity. There are two basic goals of drug treatment for lupus. The first goal is to reduce inflammation within the affected tissues. The second goal is to identify and suppress the specific abnormalities of the immune system that are considered responsible for tissue inflammation. The overall therapeutic plan generally groups lupus manifestations into four broad categories based on primary treatment modality used for initial treatment.

Fever, joint pain (arthralgias), arthritis, and serositis (pleurisy or pericarditis) can often be managed effectively by the administration of nonsteroidal, anti-inflammatory drugs (NSAIDs), such as aspirin, salicylates, ibuprofen, naproxen, clinoril, oxaprozin and tolmetin. The most common side-effects include gastrointestinal complaints and the potentiation of peptic ulcers. Acetaminophen derivatives can be safely taken with NSAIDs for added pain relief.

Cutaneous features of systemic lupus are usually most effectively managed with antimalarial drugs, such as hydroxychloroquine, chloroquine and quinacrine. Due to the high rate of cutaneous disease relapse and the safety of low-dose therapy, antimalarial

drug is usually prescribed on an indefinite basis for patients displaying lupus skin conditions. Retinoids such as isotretinoin (Acutane™) and etretinate (Acitretin™) demonstrate beneficial results when given orally, with reduction of lesions refractory to traditional antimalarial drug therapy.

5

More serious organ involvement is generally treated by the administration of a corticosteroid, given orally or intravenously. Prednisone is the most commonly used oral corticosteroid. When oral administration of steroids proves ineffective, intravenous methyl prednisolone pulse therapy (high dose) is often used in the treatment of lupus nephritis and other serious non-renal manifestations, such as hemolytic anemia, central nervous system inflammation (cerebritis), life-threatening low-platelet counts, and severe pleuropericarditis. Mild androgenic compounds such as Danazol™ and dehydroepiandrosterone (DHEA) have also been used in controlling immune thrombocytopenia and severe hemolytic anemia.

10
15

DHEA has also been reported to be effective for treating various manifestations of systemic lupus, with a focus upon the reported ability of DHEA to affect the hormonal and immune systems.

20

Immunosuppressive drugs are a fourth group used in treatment of systemic lupus. These drugs are employed when corticosteroid therapy is ineffective or intolerable for the patient. Immunosuppressive drugs include azathioprine (Imuran™), cyclosporin A (Sandimmune™), alkylating agents (nitrogen mustards, cyclophosphamide, and chlorambucil), and methotrexate. These drugs are sometimes used in combination with corticosteroids.

25

The above described treatment regimens for lupus often meet with limited success. Hence, the search continues for alternative treatments for lupus.

30

SUMMARY OF THE INVENTION

The invention is directed to the prophylactic, ameliorative and curative treatment of lupus erythematosus by administering $\Delta 5$ -androstene- 3β -ol-7,17-dione and precursors thereof which are readily metabolized *in vivo* to $\Delta 5$ -androstene- 3β -ol-7,17-dione but essentially incapable of being metabolized to androgens, estrogens or dehydroepiandrosterone.

DETAILED DESCRIPTION OF THE INVENTION INCLUDING A BEST MODE

Lupus erythematosus can be treated by administering therapeutic amounts of $\Delta 5$ -androstene- 3β -ol-7,17 dione and precursors thereof which are readily metabolized *in vivo* to $\Delta 5$ -androstene- 3β -ol-7,17-dione but essentially incapable of being metabolized to androgens, estrogens or dehydroepiandrosterone, such as $\Delta 5$ -androstene- 3β -acetoxy-7,17 dione and other 3β esters thereof.

Such treatment can be prophylactic, ameliorative or curative in nature.

Without intending to be bound by any theory, it is believed that $\Delta 5$ -androstene- 3β -ol-7,17 dione is effective for preventing, ameliorating and curing lupus erythematosus by modulating appropriate aspects of the aberrantly functioning immune system responsible for the onset and/or continuation of lupus erythematosus (*i.e.*, increasing depressed and/or decreasing hyperactive aspects of the system), particularly with respect to the production of appropriate concentrations of IL-2.

$\Delta 5$ -Androstene- 3β -ol-7,17-dione

$\Delta 5$ -androstene- 3β -ol-7,17 dione is a derivative of dehydroepiandrosterone (DHEA) which does not appreciably stimulate, increase or otherwise enhance the production of sex hormones. $\Delta 5$ -androstene- 3β -ol-7,17 dione is commercially

available from a number of sources including Steraloids, Inc. of Wilton, New Hampshire. A number of procedures are available for synthesizing Δ^5 -androstene- 3β -ol-7,17 dione from DHEA, with one such procedure described in United States Patent No. 5,296,481.

5

Precursors of Δ^5 -androstene- 3β -ol-7,17-dione, which are essentially incapable of being metabolized to androgens, estrogens or dehydroepiandrosterone, may also be usefully employed in the treatment of lupus erythematosus. Such precursors are readily metabolized *in vivo* to the active Δ^5 -androstene- 3β -ol-7,17-dione. One example of such a metabolizable precursor is the commercially available Δ^5 -androstene- 3β -acetyl-7,17-dione. The 3β -acetyl group is readily hydrolyzed *in vivo* by esterases located in the blood and various body tissue to produce the active Δ^5 -androstene- 3β -ol-7,17-dione, and is believed to be less susceptible to oxidation at the 3-position during the manufacturing process relative to Δ^5 -androstene- 3β -ol-7,17-dione.

15

Administration

20

Administration Route

The Δ^5 Androstene- 3β -acetoxy-7,17-dione can be administered by virtually any of the commonly accepted practices for the administration of pharmaceutical preparations including specifically, but not exclusively, intravenous injection, mucosal administration, oral consumption, ocular administration, subcutaneous injection, transdermal administration, etc.

25

Mucosal administration of Δ^5 Androstene- 3β -acetoxy-7,17-dione includes such routes as buccal, endotracheal, inhalation, nasal, pharyngeal, rectal, sublingual, vaginal, etc. For administration through the buccal / inhalation / sublingual / pharyngeal / endotracheal mucosa, the steroid may be formulated as an emulsion, gum, lozenge, spray, tablet or an inclusion complex such as cyclodextrin inclusion

30

complexes. Nasal administration is conveniently conducted through the use of a sniffing power or nasal spray. For rectal and vaginal administration the steroid may be formulated as a cream, douch, enema or suppository.

5 Oral consumption of the steroid may be effected by incorporating the steroid into a food or drink, or formulating the steroid into a chewable or swallowable tablet or capsule.

10 Ocular administration may be effected by incorporating the steroid into a solution or suspension adapted for ocular application such as drops or sprays.

Intravenous and subcutaneous administration involves incorporating the steroid into a pharmaceutically acceptable and injectable carrier.

15 For transdermal administration, the steroid may be conveniently incorporated into a lipophilic carrier and formulated as a topical creme or in an adhesive patch.

Dose Rate

20 The range of dosages and dose rates effective for achieving the desired biological properties and characteristics may be determined in accordance with standard industry practices. These ranges can be expected to differ depending upon whether the desired response is the prophylactic, ameliorative or curative treatment of lupus erythematosus, the type of lupus and the severity of symptoms.

25

EXPERIMENTAL

Experiment 1

(Preparation of Δ^5 Androstene-3 β -acetoxy-7,17-dione)

5

Step One:

(Preparation of Δ^5 Androstene-3-acetoxy-17-one)

10 A suitable, three-necked, round-bottom flask equipped with an overhead stirrer, reflux condenser, solids addition funnel and 110-volt temperature controller was charged with a mixture of dichloromethane (90 ml), glacial acetic acid (150 ml), and acetic anhydride (250 ml). To the mixture was added dehydroepiandrosterone (0.20 moles) purchased from Steraloids, Inc. of Wilton, New Hampshire. The mixture was stirred to dissolve the solid dehydroepiandrosterone, and anhydrous sodium acetate (35.0 g) added. The resulting mixture was heated at 75°C with stirring for 3 hours to complete the reaction.

20 The reaction mixture was poured into one liter of water and the resulting slurry stirred at room temperature for 2 hours. The organic dichloromethane layer was separated from the aqueous layer, and the aqueous layer extracted once with 50 ml of fresh dichloromethane. The combined organic dichloromethane extract was washed with water, saturated sodium bicarbonate solution (until neutral), and water. The resulting washed organic dichloromethane extract was evaporated under reduced pressure to a volume of 40 ml. Methanol (100 ml) was added to this concentrated extract and the resulting solid mass was cooled at 0°C in a refrigerator for 2 hours.

30 The resulting solid white product was collected by vacuum filtration on a Buchner funnel and the filter cake air dried on the funnel to form a first crop of product weighing 50.5 g. The filtrate mother liquor was concentrated by evaporation under reduced pressure, and cooled at 0°C in a refrigerator. The resulting solid white product was collected by vacuum filtration on a Buchner funnel and the filter cake air dried on the funnel to form a second crop of product weighing 10.2 g. The filtrate mother liquor

from the second crop of product was diluted with water and the mixture was cooled at 0°C in a refrigerator. The resulting solid white product was collected by vacuum filtration on a Buchner funnel and the filter cake air dried on the funnel to form a third crop of product weighing 4.2 g.

5

The first, second and third crops of product were combined to produce a total of 64.9 grams of Δ^5 androstene-3-acetoxy-17-one.

10

Theoretical yield	=	66.1 g
First crop yield	=	50.5 g (76.4%)
Second crop yield	=	10.2 g (15.4%)
Third crop yield	=	4.2 g (6.4%)

Step Two:

15

(Preparation of Crude Δ^5 Androstene-3-acetoxy-7,17-dione)

20

A suitable, three-necked, round-bottom flask equipped with an overhead stirrer, reflux condenser, addition funnel, thermometer, mineral oil filled bubbler and a gas inlet tube connected to a nitrogen cylinder, was charged with acetone (3.5 L) and cyclohexane (3.5 L). 1.51 moles of the Δ^5 Androstene-3-acetoxy-17-one prepared in Step One was added to the flask with stirring to dissolve the solid Δ^5 Androstene-3-acetoxy-17-one. 2.48 moles of solid sodium metaperiodate and water (1.1 L) were added to the stirred solution. 14.75 moles of a 70% aqueous solution of *t*-butyl hydroperoxide (2.0 L) was added to the flask through the addition funnel over a one-half hour period.

25

30

Over the first hour, the reaction mixture temperature rose from 20°C to 32°C. Tap water was added to an external cooling bath and the reaction mixture temperature returned to 20°C. The reaction mixture was constantly vigorously stirred throughout the experiment, and the reaction judged to be complete after 48 hours by TLC monitoring of the disappearance of starting material. The mixture changed from a white slurry to a light yellow slurry over the course of the reaction.

The reaction mixture was poured into a stirred ice/water mixture (12 kg ice and 8 L water). Potassium sulfite (3 L of a 45% aqueous solution) was then added to the diluted reaction mixture over 30 minutes (100mL/min) to destroy any remaining oxidant. The diluted mixture was stirred for an additional 2 hours, with ice added as
5 needed to maintain the mixture at 15°C.

The resulting diluted, cooled reaction mixture was transferred to a suitable container and ethyl acetate (3 L) was added to dissolve and extract the product. The resultant mixture was stirred for one-half hour and then allowed to stand so as to permit
10 the organic and aqueous layers to separate. The aqueous layer was examined by TLC, found to contain no product, and discarded. The solids containing organic layer was transferred to a separatory funnel, washed with water (3 x 1.5 L), then washed with a saturated salt solution (1 x 1.5 L). The washed organic layer was dried over sodium sulfate (300 g), with decolorizing carbon (100 g) added. The resulting organic slurry
15 was filtered through a ceramic Buchner funnel containing a 0.5 inch Celite pad (100 g). The filter cake was washed with ethyl acetate (2 x 150 ml) and the washing combined with the filtrate.

The combined organic filtrate was concentrated *in vacuo* to near dryness to
20 produce an off-white semi-solid. The semi-solid was suspended in methanol (400 ml) and again concentrated *in vacuo* to near dryness to produce a semi-solid. The semi-solid was slurried in methanol (600 ml) and the slurry stirred for 2 hours at ambient temperature. The solid product was collected by filtration on a ceramic Buchner funnel, and the solids washed with cold (5°C) methanol (2 x 75 ml). The solid
25 product was dried at 65°C for 48 hours under high vacuum (<1 mm Hg vacuum). The process yielded 232 grams of crude solid Δ^5 Androstene-3-acetoxy-7,17-dione.

Theoretical yield	=	521 g
Actual yield	=	232 g (44.5%)

30

Step Three:

(Preparation of Purified Δ^5 Androstene-3-acetoxy-7,17-dione)

METHOD A:

5

A 500 ml round bottom flask equipped with a magnetic stirrer was charged with 25.0 grams of the crude Δ^5 Androstene-3-acetoxy-7,17-dione prepared in Step Two and 300 ml of a mixture of methanol and ethyl acetate (1:1, v/v). The magnetic stirrer was activated and the slurry stirred at room temperature until the crude Δ^5 Androstene-3-acetoxy-7,17-dione was completely dissolved in the solvent mixture to form a first solution. A freshly prepared 10% aqueous solution of sodium bicarbonate (25 ml) was added over 10 minutes to the reaction mixture. The resulting milky mass was stirred at room temperature for 2.5 hours.

15

The reaction mixture was concentrated at room temperature under reduced pressure to 100 ml volume. The concentrated reaction mixture was diluted with 200 ml of ice water and stirred for 30 minutes at 0-5°C. The precipitated solids were collected on a ceramic Buchner funnel, and the aqueous filtrate reserved for additional product recovery. The solids on the funnel were washed with water (until neutral), and methanol (2 x 30 ml), with the methanol washing also reserved for product recovery. The first crop of solids was dried overnight under vacuum to give 18.0 g of purified product.

20

The aqueous filtrate from the first crop of solids was extracted with ethyl acetate (100 ml), and the separated organic extract was washed with water. The solvent of the washed extract was removed under reduced pressure to produce a solid product. This solid product was dissolved in the methanol washing from the first crop of solids, and the solution concentrated to 30 ml volume. Upon cooling the concentrate, a solid precipitate product formed which was collected by vacuum filtration. The second crop of solids was air dried to give 5.2 g of purified product.

25

30

The mother liquor filtrate from the second crop of solids was diluted with water and cooled. The resulting white solid precipitate was collected by vacuum filtration, and dried overnight at room temperature to give a third crop of 1.0 g of purified product.

5 The process yielded a total of 24.2 grams of purified solid Δ^5 Androstene-3-acetoxy-7,17-dione.

Theoretical recovery = 25.0 g

Actual recovery = 24.2 g (96.8%)

10

METHOD B:

A suitable round bottom flask equipped with a magnetic stirrer was charged with 1.0 gram of the crude Δ^5 Androstene-3-acetoxy-7,17-dione prepared in Step Two and 10 ml of acetone. The magnetic stirrer was activated and the slurry was stirred at room temperature until the crude Δ^5 Androstene-3-acetoxy-7,17-dione was completely dissolved in the acetone. To this solution was added 2.0 g of aluminum oxide basic. The resulting slurry was stirred at room temperature for 1 hour, then filtered through a bed of Celite. The collected solids and Celite bed were washed once with 5 ml of acetone, and the washing combined with the filtrate. The combined filtrate was 20 evaporated to near dryness under reduced pressure to produce a solid product. The solid product was dissolved in a mixture of methanol and isopropyl ether (8:2, v/v) with heating. This solution was cooled at 0-5°C for 30 minutes, resulting in precipitation of a white product. The precipitated solid was collected by vacuum filtration and air dried to 25 give 0.9 grams of purified solid Δ^5 androstene-3-acetoxy-7,17-dione.

Theoretical recovery = 1.0 g

Actual recovery = 0.9 g (90.0%)

I claim:

1. A treatment method comprising treating a patient in need of such treatment with an effective autoimmune disorder ameliorative amount of a steroid selected from Δ^5 -androstene- 3β -ol-7,17-dione and metabolizable precursors thereof incapable of being appreciably metabolized to androgens, estrogens or dehydroepiandrosterone.
2. The treatment method of claim 1 wherein the method comprises treating a patient with an effective autoimmune disorder ameliorative amount of a steroid selected from Δ^5 -androstene- 3β -ol-7,17-dione and Δ^5 -androstene- 3β -acetoxy-7,17-dione.
3. The treatment method of claim 1 or 2 wherein the method of treating a patient comprises treating a lupus erythematosus-affected human patient.
4. The treatment method of claim 3 wherein the method of treating a lupus erythematosus affected human patient comprises treating a human patient affected by systemic lupus erythematosus.
5. The treatment method of claim 3 wherein the method of treating a lupus erythematosus affected human patient comprises treating a human patient afflicted with lupus erythematosus-related tissue inflammation.
6. The treatment method of claim 3 wherein the method of treating a lupus erythematosus affected human patient in need of such treatment comprises treating a patient diagnosed with lupus erythematosus.
7. The treatment method of claim 4 wherein the method of treating a systemic lupus erythematosus affected human patient comprises treating a human patient afflicted with systemic lupus erythematosus.

8. The treatment method of claim 4 wherein the method of treating a systemic lupus erythematosus affected human patient comprises treating a human patient diagnosed with systemic lupus erythematosus.
9. A treatment method comprising administering an effective preventative amount of a steroid selected from Δ^5 -androstene- 3β -ol-7,17-dione and metabolizable precursors thereof incapable of being appreciably metabolized to androgens, estrogens or dehydroepiandrosterone, to an animal susceptible to lupus erythematosus.
10. The treatment method of claim 9 wherein the method of administering an effective preventative amount of a steroid comprises administering an effective preventative amount of a steroid selected from Δ^5 -androstene- 3β -ol-7,17-dione and Δ^5 -androstene- 3β -acetoxy-7,17-dione.
11. The treatment method of claim 9 or 10 wherein the method of administering an effective preventative amount of a steroid to an animal susceptible to lupus erythematosus comprises administering the steroid to a human susceptible to lupus erythematosus.
12. The treatment method of claim 11 wherein the method of administering an effective preventative amount of a steroid to a human susceptible to lupus erythematosus comprises administering the steroid to a human substantially free of lupus erythematosus-related tissue inflammation but previously afflicted with lupus erythematosus -related tissue inflammation.
13. The treatment method of claim 11 wherein the method of administering an effective preventative amount of a steroid to a human susceptible to lupus erythematosus comprises administering the steroid to a human substantially free of lupus erythematosus-related tissue inflammation but previously diagnosed with lupus erythematosus -related tissue inflammation.

14. A treatment method comprising treating a patient in need of such treatment with an effective autoimmune disorder curative amount of a steroid selected from Δ^5 -androstene- 3β -ol-7,17-dione and metabolizable precursors thereof incapable of being appreciably metabolized to androgens, estrogens or dehydroepiandrosterone.
15. The treatment method of claim 14 wherein the method comprises treating a patient with an effective autoimmune disorder curative amount of a steroid selected from Δ^5 -androstene- 3β -ol-7,17-dione and Δ^5 -androstene- 3β -acetoxy-7,17-dione.
16. The treatment method of claim 14 or 15 wherein the method of treating a patient comprises treating a lupus erythematosus-affected human patient.
17. The treatment method of claim 16 wherein the method of treating a lupus erythematosus affected human patient comprises treating a human patient affected by systemic lupus erythematosus.
18. The treatment method of claim 16 wherein the method of treating a lupus erythematosus affected human patient comprises treating a human patient afflicted with lupus erythematosus-related tissue inflammation.
19. The treatment method of claim 14 or 15 wherein the method of treating a patient in need of such treatment comprises treating a human patient diagnosed with lupus erythematosus.
20. The treatment method of claim 14 or 15 wherein the method of treating a patient comprises treating a human patient afflicted with lupus erythematosus.
21. The treatment method of claim 14 or 15 wherein the method of treating a patient comprises treating a human patient diagnosed with systemic lupus erythematosus.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/23386

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/33, 31/56

US CL :514/169, 177, 178, 179, 182

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/169, 177, 178, 179, 182

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Database WPIDS, London: Derwent Publications Ltd., AN 94-065380, WO9403176 A1, HUMANETICS CORP, "Treatment of Alzheimer's disease and immune deficiency disorders - by administration of delta-5-androsten-3beta-ol-17-one having a 7-hydroxy or oxo-substituent", abstract, 17 February 1994, Volume 08, see the entire abstract.	1-21
Y	Database WPIDS, London: Derwent Publications Ltd., AN 94-150921, WO9408589, UNIV. LELAND STANFORD JUNIOR, "Treating systemic lupus erythematosus with dehydro epiandrosterone- partic. to control proteinuria, allowing redn. in doses of corticosteroid(s)", abstract, 28 April 1994, Volume 18, see the entire abstract.	1-21

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*&*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

29 DECEMBER 1998

Date of mailing of the international search report

16 FEB 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

FREDERICK KRASS

Telephone No. (703) 308-1235